

## Determination of the prevalence of viral aetiology of diarrhea in children less than 5 years of age in Baghdad province

Basim M. Hussan\*

### الخلاصة

اجريت الدراسة تحديد مدى انتشار المسببات المرضية الفيروسية والطفيلية والجرثومية المؤدية إلى التهاب المعدة والأمعاء في الأطفال بعمر أقل من 5 سنوات مع أعراض الإسهال في محافظة بغداد من مستشفيات مختلفة.

تم جمع ما مجموعه 250 عينة براز من الأطفال من مستشفيات مختلفة من محافظة بغداد تم التحري عن الممرضات المعوية enteropathogens البكتيرية والطفيلية باستخدام الفحص المجهرى. وقد تم زرع عينات البراز على مختلف الأوساط الزرعية MacConkey, XLD, DCA, TCBS and campylobacter media, TCBS and selenite F broth وتمت تشخيص البكتيريا بواسطة المقاييس المعتمدة على (Api20E) والأموال المضادة بالإضافة إلى خصائص المستعمرات البكتيرية. واستخدمت طريقة فحص مقاييس ألاماز المناعي للأنزيم المرتبط ELISA لتحري فيروس الروتا Rotavirus ، Astrovirus و adenovirus المعوية.

تم فحص ما مجموعه 250 عينة براز الإسهال من الأطفال دون سن 5 سنوات من العمر و من هؤلاء تم تشخيص 106 طفلاً (42.4%) كانت لديهم المسببات الفيروسية وان فيروس الروتا سجل معظم الحالات 91 (36.4%)، ويأتي بعده adenovirus المعوية في 15 (6%) من المرضى.

وكان 63 (25.2%) مسببات بكتيرية منها 36 (14.4%) كانت القولونية (ETEC) و 12 (4.8%) كانت الشيغيلا الفلكسنرية، 6 (2.4%) كانت *Campylobacter jejuni* و 9 (3.6%) كانت السالمونيلا الملهبة. وكانت المتحولة الحالة للنسج 22 (8.8%)، والجياردية المبلية 22 (8.8%) الطفيليات الأكثر شيوعاً التي تم عزلها من عينات البراز

أيضاً من إجمالي 250 عينة كانت 40 عينة (16%) عينات البراز زحاري (الإسهال الدموي) ومسببات الأمراض الأكثر شيوعاً المعزولة من عينات البراز الزحاري كانت المتحولة الحالة للنسج 22 (8.8%) و 12 (4.8%) كانت الشيغيلا الفلكسنرية و 6 (2.4%) كانت *Campylobacter jejuni*

كشفت الدراسة الحالية إلى ارتفاع معدل انتشار فيروس الروتا وكان السبب الأكثر شيوعاً للإسهال في المستشفيات العراقية في الأطفال بعمر أقل من 5 سنوات .

### Abstract

Determination the prevalence of viral, parasitic and bacterial etiology of gastroenteritis in children less than 5 years of age with diarrhea in various hospitals of Baghdad province.

\*Health and medical technical college

A total of 250 stool samples were collected from various hospitals of Baghdad province. Samples were investigated for bacterial and parasitic enteropathogens using microscopic examination. Bacterial culture of stool samples was done in different culture media including MacConkey, XLD, DCA, TCBS and campylobacter media, TCBS and selenite F broth and bacterial growth was examined by analytical profile index (Api20E) system, commercial antisera and colonies characteristics. Enzyme immunoassays were used to identify Group A rotavirus, Astrovirus and enteric adenovirus.

Out of 250, 106 children (42.4%) were diagnosed as viral infection of which Rotavirus was found in the majority of cases 91 (36.4%), Adenovirus was detected in 15(6%) patients.

bacterial causes were 63 (25.2%) of which 36 (14.4%) were *Escherichia coli* (ETEC), 12 (4.8%) were *Shigella flexneri*, 6(2.4%) were *Campylobacter jejunii* and 9(3.6) were *Salmonella enteritidis*. *Entamoeba histolytica* 22 (8.8%) and Giardia lamblia 22 (8.8%) were the most frequent parasite which isolated from all diarrheal stool specimens.

Also from total of 250 samples, 40 samples (16%) were dysenteric stool specimens (bloody diarrhea) and the most common pathogens isolated from dysenteric stool specimens were *E.histolytica* 22(8.8%), *Shigella flexneri* 12(4.8%) and *Campylobacter jejunii* 6 (2.4%).

The current study revealed a high prevalence of rotavirus as most common cause of diarrhoea in children less than 5 years of age.

## Introduction

Diarrheal disease is a major public health problem throughout the world, with over two million deaths occurring each year, mostly children under 5 years of age in developing countries (1). There is a wide range of recognized enteric pathogens such as viruses, bacteria, and parasites that cause diarrhea. Among bacteria, *Shigella spp.* and diarrheagenic *Escherichia coli* (DEC) are the

most common causes of diarrheal diseases in developing countries (2).

Gastroenteritis is defined as an inflammation of the stomach, large and small intestines (3). It is one of the most common illnesses in humans worldwide (4). Although it can affect individuals of any age, it presents a significant health risk to those at extremes of age, the very young and the very old (5). It is the second most common cause of death among adults, and the leading cause of childhood death worldwide (6). In children under 5 years, there are more than 700 million cases of gastroenteritis every year. The annual mortality associated with gastroenteritis has been estimated to be 3.5 to 5 million, with the majority of deaths occurring in developing countries (7).

Viral gastroenteritis is diseases worldwide and causing significant morbidity and mortality in children.(8) Four major viral pathogens are associated with gastroenteritis, three of them are RNA viruses (Rotavirus, Norovirus, and Astrovirus) and one DNA virus (enteric Adenovirus).(9)

The aim of this study was to determine the prevalence of bacterial, parasitic and viral etiology of gastroenteritis in children less than 5 years of age presenting with diarrhoea in various hospitals of Baghdad province

### **Patients and Methods**

Study was carried out on 250 patients were suffering from diarrhea their ages below 5 years old (135 males and 115 females) attending to Baghdad Teaching Hospital, teaching laboratories ,Pediatric Al- Mansour teaching hospital in Baghdad Medicine city and Alsader hospital in Baghdad province. Stool samples were collected from each child in disposable clean containers and processed within 2 hours of collection.

Macroscopic examination of stool samples was done for the presence of worms (*Ascaris lumbricoides*, *Strongyloides stercoralis*, *Enterobius vermicularis*, *Trichuris trichiura* and *Taenia saginata*), blood and physical characteristics such as color, appearance and odor .Microscopic examination of stool samples

was done under 10x and 40x power after being stained with iodine for the presence of parasitic eggs and larvae. Bacterial culture of stool samples was done for the presence of a common enteric bacterial pathogens such as: *Salmonella*, *Shigella*, *Campylobacter*, Enteropathogenic *E.coli*, Enterotoxigenic *E.coli* and *Vibrio cholera*. These were performed by culturing the stool samples in different culture media including MacConkey, XLD, DCA, TCBS and selenite F MacConkey, XLD, DCA, campylobacter media, TCBS and selenite F broth culture media were incubated at 37°C for 24 hours and samples incubated in selenite F broth were re-cultured in XLD media for another 24 hours at 37°C. Campylobacter media were cultured at 42°C for 48 hours Any bacterial growth was examined by standard microbiological methods such as specific biochemical tests analytical profile index (Api20E) system and commercial antisera in addition to bacterial colonies characteristics.

Stools were screened for enterohemorrhagic *E. coli* by plating on sorbitol-MacConkey agar. All non-sorbitol-fermenting colonies were tested in an agglutinating assay with O157 and H7 antisera (2).

Samples were also tested for the presence of gastroenteritis viruses (Rotavirus, Adenovirus type 40/41 and Astrovirus) using antigen detection ELISA kits [ACON laboratories, incorporated (Inc), USA] as described in manufacturers instruction. In brief, washing buffer was diluted 1:10 with distilled water and samples were also diluted 1:11 with sample diluents. Hundred microliters of samples, positive and negative control were added to corresponding microtitre plate wells followed by the addition of 100Tl conjugate. Plates were then incubated at room temperature for 60 minutes before being washed 5 times with 300Tl diluted wash buffer using an ELISA washer (ETL testing laboratories Inc. USA). Hundred microliters of substrate were then added to all wells, before incubating the plates in dark for 15 minutes. Fifty microliters of stopping solution were then added before reading the results at an ELISA plate reader [E.ma, precision, microplate reader, USA ] at 450 nm.

## Results


Out of 250 children under 5 years of age were examined 106 (42.4%) were diagnosed with viral etiology of which Rotavirus was found in the majority of cases 91 (36.4%), Adenovirus was detected in 15(6%) patients.

bacterial origin were 63 (25.2%) of which 36 (14.4%) were *Escherichia coli* (ETEC), and the other bacterial enteropathogens isolated from diarrheal stool specimens were *Shigella flexneri* 12 (4.8%) and *Campylobacter jejunii* 6(2.4%) and *Salmonella enteritidis* 9(3.6). The result of the serotyping of the bacteria in the Central Public Health Laboratories/ Ministry of Health in Baghdad city has ensured that these bacteria are *Escherichia coli* (ETEC), *Shigella flexneri*, *Salmonella enteritidis* and *Campylobacter jejunii*.

Also Data illustrated in table and figure (-1) also demonstrated that the most frequent parasite which isolated from diarrheal stool specimens were *Entamoeba histolytica* (22,8.8%) and *Giardia lamblia* (22,8.8%) also data reported in table and figure (-1) showed that (37,14.8%) children were with no causative agent isolated from their diarrheal stool samples.

**Table (-1) Distribution of etiological agents which caused diarrheal stools of children under five years age**

Etiological Agent	Number of cases	Percentage of cases
Rotavirus	91	36.4%
Enteri Adeno virus 40/41	15	6%
Enterotoxigenic <i>Escherichia coli</i> (ETEC)	36	14.4%
<i>Shigella flexneri</i>	12	4.8%
<i>Campylobacter jejunii</i>	6	2.4%
<i>Salmonella enteritidis</i>	9	3.6%
<i>Entamoeba histolytica</i>	22	8.8%
<i>Giardia lamblia</i>	22	8.8%
No pathogen identified	37	14.8%
Total	250	100%



**Figure (-1) Distribution of etiological agents which caused diarrheal stools of children under five years age**

Table (-2) showed the relation between the age of child and the incidence for each one of the causative agent of diarrhea , where all children who had Rotavirus as causative agent of diarrhea were below age 3 years , in contrast all children who had *Entamoeba histolytica* or *Giardia lamblia* as causative agent of diarrhea were above two years.

Also from the same table and figure all children who have unknown causative agent of diarrhea were under two years.

**Table (-2) : Distribution of positive stool samples according to age differences.**

Age group in years	Rotavirus	Adenovirus	ETEC	Entamoeba histolytica	Shigella flexneri	Campylobacter jejunii	Salmonella enteritidis	Giardia lamblia	No agent present
< year	20	4	5	---	---	---	---	---	28
1-2 year	31	6	14	---	3	---	---	---	9
2-3 years	40	5	17	4	3	1	---	---	---
3-4 years	---	---	---	6	6	5	4	5	---
4-5 years	---	---	---	12	---	---	5	17	---
Total	91	15	36	22	12	6	9	22	37

Of the total of 250 samples, 40 samples (16%) were dysenteric stool specimens (bloody diarrhea) and the most common pathogens isolated from dysenteric stool specimens were *E.histolytica* 22 (8.8%) *Shigella flexneri* 12(4.8%) and *Campylobacter jejunii* 6 (2.4%) as were observed in table (-3)

**Table (-3) Distribution of etiological agent which caused dysenteric diarrheal stool from 150 diarrheal stool specimens.**

Etiological agent	Number of cases	Percentage of cases
<u>E.histolytica</u>	22	8.8%
<u>Shigella flexneri</u>	12	4.8%
<u>Campylobacter jejunii</u>	6	2.4%
Total	40	16%

Table (-4) showed that non dysenteric stool specimens were observed in (84%) of all diarrheal stool specimens and the pathogens which isolated from non dysenteric stool were rotavirus (36.4%), adenovirus (6.0%) , Escherichia coli 14.4%, Salmonella enteritidis (4.7%) , Giardia lamblia 8.8% and unknown etiological agent (14.8%) .

**Table (-4) : Etiological Agent which caused non dysenteric diarrheal stool**

Etiological Agent	Number of cases	Percentage of cases
Rotavirus	91	36.4%
Adenovirus	15	6.0%
Enterotoxigenic <u>Escherichia coli</u> (ETEC)	36	14.4%
<u>Salmonella enteritidis</u>	9	3.6%
<u>Giardia lamblia</u>	22	8.8%
Unknown etiological agent	37	14.8%
Total	210	84%

## Discussion

Diseases causing diarrhoea are the most important cause of morbidity and mortality in developing countries. Five hundred million cases of acute diarrhoea occur annually in children aged less than five years throughout the world.(10,11)

The sex distribution among the children in our study showed a slight preponderance of diarrhoea in males compared with females (135 males and 115 females). Similar findings have been reported in other studies in Iraq (12,13,14)

In the present study, enteropathogens were detected in 213/250 (85.2%) paediatric patients under five years of age. The present study clearly shows that Rotavirus is the most common cause of acute gastroenteritis among children 90 (36.4%). This agrees with Al-Khashab (12) and Al-Kaby (13) from Iraq, as well as (15) and Kamalaratnam et al. from India (16), and Gusmao et al. from Brazil (17) however, our rotavirus prevalence is slightly lower than that reported in Central Saudi Arabia, the prevalence was 44.3% in the under 5 year age group

In a study conducted in Karachi, Pakistan, Rotavirus was detected in 112/818 (13.7%) of stool samples from infants with diarrhea (18). However, in another study carried out in India, Rotavirus was found in 22.6% of pediatric patients, who fell into the age group <1-3 years (19). In the United Kingdom, Rotavirus infection prevalence was found to be 36% in children under 16 years of age (20) while the prevalence rate of Rotavirus in different cities of South Africa varied from 13% to as high as 55% in under 5 year of age (21) and in the USA the rate was 6.8% in children less than 6 years of age (22). In Lima, Peru, rotavirus was significantly associated with acute diarrhea( AD) as compared to children with persistent diarrhea (23). In northeastern Brazil, only 3.7% of children presenting to a clinic had rotavirus identified in their stool.(24)

Adenovirus was the second most common virus to be found with a prevalence rate of 6%. Other studies done in different parts of the world, showed prevalence rates ranging from as low as 0.8% in under 6 years old in the USA(22) to as high as 28% in children

under 12 years of age in Ghana.(25) Our results were in accordance with the reported prevalence of Adenovirus but this is in contrast to one study which found Astrovirus to be more common than Adenovirus.(22) When compared to other studies done worldwide it was reported that the prevalence of Astrovirus ranged from 0.4% in the under 5 year old age group in Tanzania(26) to 9.2% in <1 to 4 year old children in Saudi Arabia (27) .

Bacteria isolated in the current study were 25.2%, of which *Escherichia coli* (ETEC) made up 14.4%, 4.8% was due to *Shigella flexneri* ,3.6% was due to *Salmonella enteritidis* and 2.4% was due *Campylobacter jejunii*. The result of the serotyping of the bacteria in the Central Public Health Laboratories/ Ministry of Health in Baghdad city has ensured that these bacteria are *Escherichia coli* (ETEC), *Shigella flexneri*, *Salmonella enteritidis* and *Campylobacter jejunii*

In Brazilian children, ETEC was found to be associated with persistent diarrhea. (28) Whereas, in rural India, ETEC was found significantly more often in children with acute diarrhea than in controls.(29) ETEC was identified more frequently in children with PD compared to AD in rural Indian, but the association was not statistically significant.(30) In Vietnam, in a sample of children experiencing PD, ETEC was the most frequently identified etiologic agent

Salmonellosis has considered one of the most common causes of diarrheal diseases in Iraq by (31) who found that children more than five years old are more susceptible to Salmonellosis and with study of (32) who record that the Children aged 2 are the most affected age group (incidence 391.6 per 100 000 population).

In two other studies done worldwide, Salmonella was found in 5.8% of patients in Australia in <1 up to 14 years of age(33) and Shigella was found in 6% in Palestine in children under 5 years of age.(34)

In addition our study's demonstrated that the most frequent parasite which isolated from diarrheal stool specimens were *Entamoeba histolytica* (22,8.8%) and *Giardia lamblia* (22,8.8%)

and all children who had *Entamoeba histolytica* or *Giardia lamblia* as causative agent of diarrhea were above 2 years age. We believe that children who aged younger than 2 years are under the supervision and care of their mothers, while older children are distance from their mothers and they play more that make them near from contaminated water sources that are the principle source of *Entamoeba histolytica* and *Giardia lamblia*. *Giardia lamblia* is a common intestinal protozoa present in 20-30% of people in the developing world.(35). In children living in northeastern Brazil, *Giardia* was frequently identified in children with persistent In other studies, an association was found between *Giardia* and PD in northeastern Brazil and in a New Delhi slum.(36) however, *Giardia* was identified infrequently in children living in Bangladesh and there was insignificant association

In this results have shown that 40 (16%) were dysenteric stool specimens (bloody diarrhea) and the most common pathogens isolated from dysenteric stool specimens were *E.histolytica* 22 (8.8%) *Shigella flexneri* 12(4.8%) and *Campylobacter jejunii* 6 (2.4%)

During the period of the current study, no clinical evidence of vibriosis due to *Vibrio cholerae* or food poisoning due to *Staphylococcus aureus* was found. other bacteria that are sometimes incriminated in diarrhoea among children, such as *Yersinia enterocolitica* (15)

This study have reported that 37(14.8%) children were with no pathogenic agent isolated from their diarrheal stool samples.

We have noticed that all children who have no pathogen identified of diarrhea were under two years, to interpret the high percentage of unknown causative agent of diarrhea ,this may be occurred following a change in the type of milk intake by child or the addition of a new type of food for the child, especially if it was not appropriate for the age also numerous other factors not related etiology have been identified as characteristics influencing of diarrhea. These include malnutrition (37-39) maternal education level(40,41) previous illness(42,43) time of weaning.

## Conclusion

The current study revealed a high prevalence of rotavirus as most common cause of diarrhoea in children less than 5 years of age

## Recommendation

Accordingly, we recommend routine general stool examination, stool culture/sensitivity and detection of rotavirus and enteri Adeno virus by ELISA methods for all patients with diarrhoea to identify the causative agent involved.

## References

1. PulseNet USA.( 2004). One day (24–28 h) standardized laboratory protocol for molecular subtyping of Escherichia coli O157:H7, nontyphoidal Salmonella serotypes, and Shigella sonnei by pulsed-field gel electrophoresis (PFGE). CDC, National Center for Infectious Diseases, Atlanta, GA.
2. Davidson, G., G. Barnes, D. Bass, M. Cohen, A. Fasano, O. Fontaine, and S. Guandalini.( 2002). Infectious diarrhea in children: working group report of the First World Congress of Pediatric Gastroenterology, Hepatology, and Nutrition. *J. Pediatr. Gastroenterol. Nutr.* 35: 2 S143–S150.
3. Jones M, Harrach B, Ganac R, Gozum M, Dela Cruz W, Riedel B.( 2007). New adenovirus species found in a patient presenting with gastroenteritis. *J Virol.* 81: 5978-84
4. Okitsu-Negishi S, Nguyen T, Phan T, Ushijima H.( 2004). Molecular epidemiology of viral gastroenteritis in Asia. *Pediatr Int.* 46: 245-52.
5. Jones S. A.( 2003). clinical pathway for pediatric gastroenteritis. *Gastroenterol Nurs.* 26: 7-18.
6. Zamir D, Weiler Z, Kogan E, Ben-Valid E, Hay E, Reitblat T. (2006). Singledose quinolone treatment in acute gastroenteritis. *J Clin Gastroenterol.* 40: 186-90.
7. Wilhelmi I, Roman E, Sánchez-Fauquier A.( 2003). Viruses causing gastroenteritis. *Clin Microbiol Infect.* 9: 247-62
8. Chen S, Chang Y, Lee Y, Chao H, Tsao K, Lin T.( 2007). Molecular epidemiology and clinical manifestations of viral gastroenteritis in hospitalized pediatric patients in Northern Taiwan. *J Clin Microbiol.* 45:

9. Nguyen TA, Yagyu F, Okame M, Phan TG, Trinh QD, Yan H. (2007). Diversity of viruses associated with acute gastroenteritis in children hospitalized with diarrhea in Ho Chi Minh City, Vietnam. *J Med Virol.* 79: 582-90.
10. Walsh A, Warren S. (1979). Selective primary health care: an interim strategy for disease control in developing countries. *N Engl J Med.* 301: 967-74.
11. Snyder D, Merson H. (1982). The magnitude of the global problem of acute diarrhoeal disease: a review of active surveillance data. *Bull World Health Organ.* 60: 605-13
12. Al-Khashab FM. (2001). Microbiological study of enteric pathogens associated with persistent diarrhea among children under five years [MSc thesis]. Tikrit, Iraq, College of Medicine, University of Tikrit.
13. Al-Kaby FJ. (2000). A study on diarrhea in relation to malnutrition in children under 2 years in Baghdad [MSc thesis]. Baghdad, Iraq, College of Medicine, Al-Mustansiriya University.
14. Al-Jebouri HS. (2001). Aetiology of diarrheal illness of children at Tikrit Teaching Hospital [MSc thesis]. Tikrit, Iraq, College of Education, University of Tikrit.
15. Horan TC, Andrus M, Dudeck MA. (2008). CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *American journal of infection control.* 36:309–32
16. Kamalaratnam CN. (2001) A prospective study of nosocomial enteric pathogen acquisition in hospitalized children in South India. *Journal of tropical pediatrics.* 47(1):46–9.
17. Gusmão RH. (1995). Rotaviruses as a cause of nosocomial, infantile diarrhoea in northern Brazil: pilot study. *Memorias do Instituto Oswaldo Cruz.* 90(6):743–9.
18. Nishio O, Matsui K, Oka T, Ushijima H, Mubina A, Dure-Samin A. (2000). Rotavirus infection among infants with diarrhea in Pakistan. *Pediatr Int.* 42: 425-7.
19. Saravanan P, Ananthan S, Ananthasubramanian M. (2004). Rotavirus infection among infants and young children in Chennai, South India. *Indian J Med Microbiol.* 22: 212-21.
20. Cunliffe N, Allan C, Lowe S, Sopwith W, Booth A, Nakagomi O. (2007). Healthcare-associated rotavirus gastroenteritis in a large paediatric hospital in the UK. *J Hosp Infect.* 67: 240-4.

21. Steele A, Peenze I, de Beer M, Pager C, Yeats J, Potgieter N.( 2003). Anticipating rotavirus vaccines: epidemiology and surveillance of rotavirus in South Africa. *Vaccine*. 21: 354-60.
22. Rodriguez-Baez N, O'Brien R, Qiu SQ, Bass DM. (2002). Astrovirus, Adenovirus, and Rotavirus in Hospitalized Children: Prevalence and Association With Gastroenteritis. *J Paediat Gastroenterol Nutrition*. 35: 64.
23. Lanata CF, Black RE, Maurtua D.( 1992). Etiologic agents in acute vs persistent diarrhea in children under three years of age in peri-urban Lima, Peru. *Acta Paediatr Suppl*.381:32-8
24. Lima AA, Fang G, Schorling JB.( 1992).Persistent diarrhea in Northeast Brazil: etiologies and interactions with malnutrition. *Acta Paediatr*.81 Suppl 381:39-44.
25. Reither K, Ignatius R, Weitzel T, Seidu-Korkor A, Anyidoho L, Saad E. (2007).Acute childhood diarrhoea in northern Ghana: epidemiological, clinical and microbiological characteristics. *BMC Infect Dis*. 7: 104
26. Moyo S, Gro N, Kirsti V, Matee M, Kitundu J, Maselle S.( 2007). Prevalence of enteropathogenic viruses and molecular characterization of group A rotavirus among children with diarrhea in Dar es Salaam Tanzania. *BMC Public Health*. 7: 359.
27. Meqdam M, Thwiny I . (2007).Prevalence of group a rotavirus, enteric adenovirus, norovirus and astrovirus infections among children with acute gastroenteritis in Al-Qassim, Saudi Arabia. *Pak J Med Sci*. 23: 551-5.
28. Lima AA, Moore SR, Barboza MS, Jr., et al.( 2000). Persistent diarrhea signals a critical period of increased diarrhea burdens and nutritional shortfalls: a prospective cohort study among children in northeastern Brazil. *J Infect Dis*.181:1643-51.
29. Bhan MK, Sazawal S, Raj P. (1989).Aggregative *Escherichia coli*, *Salmonella*, and *Shigella* are associated with increasing duration of diarrhea. *Indian J Pediatr*.56:81-6.
30. Bhan MK, Bhandari N, Sazawal S. (1989). Descriptive epidemiology of persistent diarrhoea among young children in rural northern India. *Bull World Health. Organ* 67:281-8.
31. Al-Hayali, G.M.F. (1993). Salmonellosis in Iraq, A five years review 1985- 1989. Diploma thesis, College of Medicine / AL- Nahrian.

32. Czerwinski, M.; Czarkowski, M.P. and Baumann, A. (2008). Salmonellosis in Poland in 2006. *Przegl. Epidemiol.* 62(2): 301-310.
33. Saravanan P, Ananthan S, Ananthasubramanian M. Rotavirus infection among infants and young children in Chennai, South India. *Indian J Med Microbiol* 2004. 22: 212-21
34. Abu-Elamreen F, Abed A, Sharif F. *Viral.* (2008).bacterial and parasitic etiology of pediatric diarrhea in Gaza, Palestine. *Med Princ Pract.*17: 296-301.
35. Floch MH. (2010).*Giardia lamblia* and Other Protozoan Infections. In: Floch MH, Floch, N.R., Kowdley, K.V., Pitchumoni, C.S., Rosenthal,, R.J., Scoplapio, J.S., ed. *Netter's Gastroenterology.* 2nd ed: Saunders Elsevier.
- Lima AA, Moore SR, Barboza MS, Jr.( 2000). Persistent diarrhea signals a critical period of increased diarrhea burdens and nutritional shortfalls: a prospective cohort study among children in northeastern Brazil. *J Infect Dis.*181:1643-51
36. Bhandari N, Bahl R, Dua T, Kumar R, Srivastava R.( 1999). Role of protozoa as risk factors for persistent diarrhea. *Indian J Pediatr.* 66:21-6.
37. Black RE. Persistent diarrhea in children of developing countries.( 1993). *Pediatr Infect Dis J.*12:751-61.
38. Thanh PN, Ly DT, Dung PT, Le PD. (1992).Clinical aspects of acute vs persistent diarrhea in Ho Chi Minh City, Vietnam. *Acta Paediatr Suppl.*381:121-3.
39. Lima AA, Guerrant RL. (1992).Persistent diarrhea in children: epidemiology, risk factors,pathophysiology, nutritional impact, and management. *Epidemiol Rev.*14:222-42
40. Fraser D, Dagan R, Porat N.( 1998). Persistent diarrhea in a cohort of Israeli Bedouin infants: role of enteric pathogens and family and environmental factors. *J Infect Dis.*178:1081-8
42. Moore SR, Lima NL, Soares AM.( 2010). Prolonged Episodes of Acute Diarrhea Reduce Growth and Increase Risk of Persistent Diarrhea in Children. *Gastroenterology.*139:1156-64.
43. Persistent diarrhoea in children in developing countries: memorandum from a WHO meeting. *Bull World Health Organ.*66:709-17.