

The use of *Omp W* Gene in Detection of *Vibrio cholerae* Isolated from Diarrhea Cases of Children in AL-Diwaniya Province.

Thuraya Aamer Habeeb, Adnan H. Al-Hamadani* and Jawad K. AL-Janabi**.

الخلاصة

تم خلال هذه الدراسة جمع 221 عينة براز من اطفال تحت سن الخامسة عشريعتون من اسهال مائي ومن كلا الجنسين والذين راجعوا مستشفى الولادة والاطفال في محافظته الديوانيه خلال الفتره الزمنيه من تشرين الاول 2008 وحتى حزيران 2009. كذلك تم جمع 27 عينة ماء من ثلاثه مناطق مختلفه لنهر الديوانيه وبنفس الفتره الزمنيه لغرض دراسه الصفات المظهرية والجزيئيه لضمات الكوليرا وذلك لغرض تقييم طرق التشخيص المختبرية الروتينية المستخدمة في تشخيص سلالات ضمات الكوليرا المختلفه ومقارنتها بتقنيه تفاعل السلسله المتبلمره (PCR).

تم عزل وتشخيص ضمات الكوليرا باستخدام طريقه الزرع اضافه الى الاختبارات الكيموحياتيه ونظام التشخيص (API 20E). كذلك اظهرت نتائج التنميط المصلي باستخدام (monovalent Ogawa و Polyvalent *V. cholerae* O1 antisera و monovalent Inaba) ان اغلب ضمات الكولير المعزوله سريريا كانت تعود الى النمط المصلي (O1) بينما جميع ضمات الكوليرا التي عزلت من المياه السطحيه لنهر الديوانيه كانت تعود الى النمط المصلي (Non- O1).

تم استخدام فحص تفاعل السلسله المتبلمره (PCR) للكشف عن جين (*ompW*) الذي يشفر لبروتين الغشاء الخارجي الخاص بضمات الكوليرا. اعتمادا على نتائج فحص تفاعل البلمره، اظهرت هذه الدراسه ان معدل عزل ضمات الكوليرا كان 5.9% بالنسبه لعينات البراز بينما 14.8% بالنسبه لعينات الماء. كذلك اظهرت نتائج فحص تفاعل السلسله المتبلمره (PCR) ان هنالك خصوصيه عاليه (100% و 100% و 97% و 86%) في الكشف عن سلالات ضمات الكوليرا ضد كل من الطرق الروتينية المختلفه كالزرع، الاختبارات الكيموحياتيه، نظام (API 20E) والتنميط المصلي على التوالي.

Abstract

A total of 221 stool samples were collected from children suffering from watery diarrhea, less than 15 years old of both genders whom admitted to the Maternity and Children Teaching Hospital in Al- Diwaniya Province, In addition to that, 27 water samples were also collected from three different loci of AL-Diwaniya River, at the period of October 2008 to June 2009,

* College of Medicine / Al-Qadisiya University- Iraq.

** Maternity and Children Teaching Hospital/Al- Diwaniya health office-Iraq.

in order to evaluate the routine laboratory diagnostic procedures in the diagnosis the multi-serogroups or serotype of *Vibrio cholerae* strains and compared them with molecular technique as Polymerase chain reaction (PCR).

Vibrio cholerae has been isolated and identified by using culturing method in addition to biochemical tests, API 20E diagnostic kit. Serotyping by using polyvalent *Vibrio cholerae*, O1 antisera, and monovalent Ogawa and Inaba revealed that the most of clinical *V. cholerae* were of serogroup O1, while all the *V. cholerae* isolated from surface water of AL- Diwanyia river was Non-O1 serogroup. PCR technique was used to detect ompW gene encoding to outer membrane protein of *V. cholerae*. Based on the PCR results, the rate of *Vibrio cholerae* isolation from stool samples was 5.9%. PCR results showed that there were high specificity (100%, 100%, 97% and 86%) in detection of *Vibrio cholerae* strains versus each of cultural, biochemical, API 20E system and serological tests, respectively.

Introduction

Cholera has been recognized as a killer disease since earliest time. The disease is caused by infection of the small intestine by *Vibrio cholerae* O1 and O1391 which is characterized by severe dehydrating diarrheal condition and is one disease in modern times that is epidemic, endemic and pandemic in nature².

Cholera is a historically-feared epidemic of diarrheal disease that remains a major public health problem in many parts of Africa, Asia, and Latin America³. Though rare in developing countries, cholera is still an important infection worldwide⁴. Also cholera has been categorized as one of the “emerging and reemerging infections”⁵ threatening many developing countries. Outbreaks of cholera cause deaths estimated at 120,000 annually worldwide and many more cases each year, of which the vast majority occurs in children⁶.

In the last decades, attention to cholera epidemiology increased, as cholera epidemics became a worldwide health problem. Detailed investigation of *V. cholerae* interactions with its host and with other organisms in the environment suggest that cholera dynamics are much more complex than previously times⁷. *V. cholerae* O1 and O139 are currently believed to be the only serogroups causing epidemic cholera⁸. The natural inhabitant of *Vibrios* is the aquatic environment and water plays an important role in disease transmission⁹.

In the past, the ability to differentiate *V. cholerae* strains for epidemiological purposes was hampered by the low discriminatory capability of available systems, such as biochemical and serologic identification¹⁰. Molecular techniques should be considered as potentially important tools in the effective monitoring of strains of *V. cholerae* especially those associated with epidemiologically well-defined outbreaks. So, application of molecular techniques to the analysis of strains has enhanced epidemiologic investigations and understanding of the sources and vehicles of transmission¹¹.

Materials and Methods

1- Samples collection

A total of 221 stool samples were obtained from patients who attended to Maternity and Children Teaching Hospital in Al-Diwanyia province, during the period from October 2008 to the June 2009. Patients mainly included children under 15 years old from both genders suffering from watery diarrhea. The stool samples were collected in sterile plain plastic containers. Then, a small amount of samples was injected in to 5 ml of alkaline peptone water (APW) prepared earlier in sterile tubes. In addition to that, 27 water samples were also collected from three different loci of AL-Diwaniya River, by using the membrane filter technique, a 500-1000 ml of water samples were collected, concentrated on 0.22 μm Pore – diameter filters, followed by incubation of the membranes in an enrichment medium of alkaline peptone water (APW) over night at 37 °C.¹².

2- Samples processing

A loopful of the culture broth, was taken from the top layer of the APW then streaked on to Thiosulphate Citrate Bile salt Sucrose agar (TCBS) and incubated overnight at 37 °C. Two-three yellow, flat, 1-3 mm diameter isolated colonies were picked from each sample and streaked on blood agar and MacConkey agar for further testing^{12, 13}.

3- Isolates identification

The isolates were further identified by microscopic examination and biochemical tests^{13, 14}. Also the isolates confirmed biochemically by using API 20-E system (Biomermex / france) according to instructions of supplied company.

4- Serotyping

Serologic confirmation was done using polyvalent, anti-Ogawa, and anti-Inaba antisera from Plasmatec Laboratory Products Ltd (Plasmatec/ UK)15.

5- Biotyping

The biotype of *V. cholerae* serogroup O1 strains was distinguished by the following methods; polymyxin B susceptibility, hemolysin test, and Voges-Proskauer test15.

6- Polymerase chain reaction (PCR) detection method

A-Method of DNA extraction: DNA extraction of isolated colony of *Vibrio cholerae* was done according to DNA extraction protocol of manufacturing company instructions (Cinagen / Iran) (Figure 1).

B- Method of PCR Reaction: The PCR amplification was performed using assay based on repetitive sequence *ompW* gene, the primers forward (5'-CAC CAA GAA GGT GAC TTT AAT TGT G-3') and revers (5'-GAA CTT ATA ACC ACC GCG -3'),16. Also the laboratory protocol was done according to manufacturing company instructions (Labaqua / Spanish).

Statistical analysis

The results were analyzed statistically by Chi-square (X^2) test at the level of significant when P -value < 0.01 . The specificity, sensitivity and diagnostic accuracy of the results were also calculated by applying the following equations: $a / a + b \times 100 = \text{sensitivity}$, $d / c + d \times 100 = \text{specificity}$.

a = Total of true positive, b = Total of false positive, c = Total of false negative and d = Total of true negative, positive and negative predictive value also calculated. Diagnostic accuracy = sensitivity + specificity / 2.17.

Results and discussion

This study shows that traditional typing systems based on serotype, biotype and other traditional methods are not always able to discriminate and identification these strains that isolated. Serotyping by using polyvalent cholera O1 antisera and monovalent Ogawa and Inaba revealed that the most of clinical *V. cholerae* were of serogroup O1, while all the *V. cholerae* isolated from surface water of AL- Diwanyia river was Non-O1 serogroup and all the

agglutinating strains in this study were identified as *V. cholerae* O1, biotype El Tor, serotype Ogawa.

This study showed that out of 221 stool samples and 27 water samples, (33) stool samples (14.9%) and (9) water samples (33.3%) were positive for culture method while 16 stool samples and 5 water samples gave positive by biochemical tests. On the other hand, 13 stool samples and 3 water samples showed positive results in API 20E diagnostic kit while only 12 stool samples gave positive result for serotyping by polyvalent *V. cholerae* O1 antisera. PCR technique showed only 13 stool samples and 4 water samples were positive.

The results of this study also revealed that all strands of DNA which resulted from the binding between specific primers and positive isolates extracted DNA appear as single band with 588 bp under the U.V light using ethidium bromide as a specific DNA stain in addition to band with the 650 pb work as Internal Positive Control (IPC) (Figure 2).

Comparison Between The PCR Method and Other Tests Used for Detection of *Vibrio cholerae*

a- PCR Results Versus Culture Method

From 48 samples were tested by culture and PCR, there were 42 samples positive by culture method versus 17 samples positive by PCR. So, the sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) and accuracy rate of PCR were (40%, 100%, 100%, 19% and 70%) respectively as demonstrated in (Table 1).

b- PCR Results Versus Biochemical Tests

From 48 samples were tested by biochemical tests and PCR, there were 21 samples positive by biochemical tests versus 17 samples positive by PCR. So the sensitivity, specificity, PPV, NPV and accuracy rate of PCR were (81%, 100%, 100%, 87% and 90.5%) respectively (Table 2).

c- PCR Results Versus API 20E

From 48 samples were tested by API 20E and PCR, there were 16 samples positive by API 20E versus 17 samples positive by PCR. The sensitivity, specificity, PPV, NPV and accuracy rate of PCR were (100%, 97%, 94%, 100% and 98.5%) respectively (Table 3).

d- PCR Results Versus Serotyping

From 48 samples were tested by API 20E and PCR, there were 12 samples positive by serotyping versus 17 samples positive by PCR. The sensitivity, specificity, PPV, NPV and accuracy rate of PCR were (100%, 86%, 70%, 100% and 93%) respectively (Table 4).

This study found that cholera infection in AL- Diwaniya province considered one of the causes of diarrhea in children of age more than one years old. This may be reflected that *Vibrio cholerae* are one of etiological agent of diarrhea that infected human and the infections occurred in a regular seasonal pattern and are particularly associated with poverty and poor sanitation.

The traditional identification of *V. cholerae* is usually achieved through a series of biochemical tests after their growth and isolation on a selective plating medium e.g., TCBS agar¹⁸. Several studies conducted in Iraq revealed that the prevalence of *Vibrio cholerae* were (42.9%), (47.14%), (87%) which reported by AL-Obidi, (2005); Kadhim,(2009); AL-Karkhy, (2005) respectively^{19, 20, 21}.

This study revealed significant differences between culture method and PCR for diagnosis of *Vibrio cholerae* and the PCR more sensitive and specific than culture method because the culture method for *Vibrio cholerae* had very poor sensitivity and numerous false positive results was occurred^{22,23}. Many bacteria can produce colonies on TCBS agar whose appearance is similar to the appearance of *V. cholerae* colonies and these false positive results take more time in their identification²⁴.

According to the reading of biochemical tests, this study showed that from 42 isolate were positive by culture only 21 isolate was positive as *Vibrio cholerae* and the other 21 isolate were non *Vibrio cholerae*. This result was different with results of API 20E diagnostic kit which found that only 16 isolates were positive as *V. cholerae*. This study showed that there were no significant differences between PCR and biochemical tests and API 20E system in the diagnosis of *Vibrio cholerae* O1. The API 20E system is indeed considered an acceptable method for the identification of the more commonly-occurring members of the family Vibrionaceae²⁵ even if there are very few reports expressly concerned with the ability of commercial systems to identify members of the genus *Vibrio*²⁶. Baron et al., (2007) showed that the direct use of the PCR-based method for the definite identification of the screened colonies gave better results than the API 20E method¹⁶. Also, West et al. (1983) mentioned that the series of biochemical tests commonly used to identify *V. cholerae* was originally designed for clinical samples²⁷.



In this study, serotyping of isolates by polyvalent O1 antisera we found 12 isolate belong to *Vibrio cholerae* O1 while the result of PCR were 17 isolate which belong to *Vibrio cholerae*. There were no significant differences between PCR and serotyping for diagnosis of *Vibrio cholerae* and we found PCR is more specific than serotyping in the diagnosis of *Vibrio cholerae* O1. Techniques based on serology are critical, but they are useful only for detecting specific serogroups. In contrast, the PCR-based identification techniques are accurate, sensitive, and permit a large throughput²⁸.

Detection methods for *Vibrio cholerae* can be divided into two categories, conventional and rapid techniques. Conventional laboratory diagnosis depends upon isolation by culture and identification by biochemical and serological tests. This process, however, is time-consuming, take about 2 days, not always accurate because close relatedness among *V. cholerae* and certain other members of the *Vibrio* spp. or *Aeromonas* spp., with respect to their biochemical properties, has often made unambiguous identification of the organism quite difficult also it may be quite expensive for a laboratory^{28, 29}. A rapid and accurate method is essential for the detection and screening of the public health threat *V. cholerae*. Hence, PCR method represents an assay format for the detection of *V. cholerae* and the DNA amplification technique is a rapid and very sensitive technique used^{22, 30, 31}.

Table (1) Validity of PCR test for diagnosis of *V. cholerae* confirmed with culture method.

Test		PCR method		Total
		+ve	-ve	
Culture method	+ve	17	25	42
	-ve	0	6	6
Total		17	31	48

Table (2) Validity of PCR test for diagnosis of *V. cholerae* confirmed with biochemical tests.

Test		PCR method		Total
		+ve	-ve	
Biochemical tests	+ve	17	4	21
	-ve	0	27	27
Total		17	31	48

Table (3) Validity of PCR test for diagnosis of *V. cholerae* confirmed with API 20E.

Test		PCR method		Total
		+ve	-ve	
API 20E	+ve	16	0	16
	-ve	1	31	32
Total		17	31	48

Table (4) Validity of PCR test for diagnosis of *V. cholerae* confirmed with serotyping.

Test		PCR method		Total
		+ve	-ve	
serotyping	+ve	12	0	12
	-ve	5	31	36
Total		17	31	48




Figure 1: DNA extraction of *Vibrio cholerae* isolates using 1% agarose gel electrophoresis.




Figure 2: DNA amplification of a 588pb fragment of Omp W gene, Line 1, 1000 bp ladder , Line 2,3,4,5 positive specimens, line 6, positive control, line 7, negative control. Amplification samples was directly loaded in a 2% agarose gel containing 0.5 mg/ml ethidium bromide with adding loading buffer in electrophoresis.

References

- 1- Rabbani, G.H. and Greenough, W.B. (1999). Food as a vehicle of transmission of cholera. *J. Diarrhoeal Dis. Res.*, 17: 1-9.
- 2- World Health Organization. (2002). Cholera in 2001. *Weekly Epidemiology Record*. 77:257-268.
- 3- Shears P. (2001). Recent developments in cholera. *Curr. Opin. Infect. Dis.* 14: 553 – 558.
- 4- Seas, C. and Gotuzzo, E. (2000). *Vibrio cholerae*. In: Mandell, G.L.; Bennett, J.E.; Dolin, R.; eds. *Principles and Practice of Infectious Diseases*. USA: Churchill Livingstone. 2266 – 2272.
- 5- Satcher, D. (1995). Emerging infections: getting ahead of the cure. *Emerg. Infect. Dis.* 1:1–6.
- 6- World Health Organization. (1995). Meeting on The Potential Role of New Cholera Vaccines in the Prevention and Control of Cholera Outbreaks during Acute Emergencies. Document CDR/GPV/95.1. World Health Organization, Geneva, Switzerland.
- 7- Codeco, C.T. (2001). Endemic and epidemic dynamics of cholera: the role of the aquatic reservoir. *BMC Infect. Dis.* 1:1.
- 8- Colwell, R. R.; Huq, A.; Chowdhury, M. A. R.; Brayton, P. R. and Xu, B. (1995). Serogroup conversion of *Vibrio cholerae*. *Can. J. Microbiol.* 41, 946–950.
- 9- Colwell, R. R.; Kaper, J. and Joseph, S. W. (1977). *Vibrio cholerae*, *Vibrio parahaemolyticus*, and other vibrios: occurrence and distribution in Chesapeake Bay. *Science*, 198, 394–396.
- 10- Mintz, E. D.; Popovic, T. and Blake, P. A. (1994). Transmission of *Vibrio cholerae* O1. In: *Vibrio cholerae* and cholera: Molecular to Global Perspectives. . Edited by Wachsmuth, K.; Blake, K. A. And Olsvik, O. American Society for Microbiology, Washington, D C. Pp.346-7.
- 11- Wachsmuth, I. K.; Evins, G. M.; Fields, P. I.; Olsvik, O.; Popovic, T.; Bopp, C. A.; Wells, J. G.; Carrello, C. and Blake, P. A.(1993). The molecular epidemiology of cholera in Latin America. *J. Infect. Dis.* 167: 621-626.
- 12- Oliver, J. D. and Kaper, J. B. (1997) *Vibrio* species. In: *Food Microbiology: Fundamentals and Frontiers*. Edited by Doyl, M. P.; Beuchat, L. R. and Montville, T. J., ASM press, Washington D C.,USA. Pp. 228-260.
- 13- Collee, J.G. ; Fraser, A.G.; Mermion, B.P. and Simmons, A. (1996). Mackie and MacCatney. *Practical Medical Microbiology*. P.425. (14th ed.). Churchill Livingstone Philadelphia.

- 14- Perilla, M.J. ;Ajello ,G; Boop ,C.; Elliott, J. ;Facklam ,R.; Popovic,T. and Wells ,J. (2003). Manual for laboratory identification and antimicrobial susceptibility testing of bacterial pathogens of public health importance in developing world, Center for disease control and prevention, Atlanta, Georgia, USA.
- 15- Elliot, E. L.; Kaysner, C. A.; Jackson, L. and Tamplin, M. L. (2001). *Vibrio cholerae*, *V. parahemolyticus*, *V. Valnificus* and other *Vibrio* spp. In: Food and Drug Administration: Bacteriological Analytical Manual, chapter 9, 8th ed. Edited by Merker, R. L., AOAC International, Gaithersburg, MD.
- 16- Baron, S.; Chevalier, S. and Lesne, J. (2007). *Vibrio cholerae* in the environment: a simple method for reliable identification of the species. *J. Health Popul. Nutr.* 25:312-318.
- 17- Niazi, A.D. (2000). Statistical Analysis in medical research. Republic of Iraq. Al- Nehrien University. P.148.
- 18- Sakazaki R. (1992) Bacteriology of *Vibrio* and related organisms,p.37-55. In D. Barua, and W. B. Greenough III (ed.),Cholera. PlenumMedical Book Company, New York, N.Y.
- 19- AL-Obidi, I.M.A. (2005). Determination of Some Epidemiological Markers in Locally Isolated *Vibrio cholerae*.Thesis,College of Science, University of AL-Mustansiryah.
- 20- Kadhim, S. A. (2009). Detection of some bacterial causes of watery diarrhea in the province of Baghdad and some northern governorates, with a study of pathological effects. M.Sc.Thesis, College of Science, University of Baghdad.
- 21- Al-Karkhy, K. A. (2005).Bacteriological and enzymatic study for somevirulence factors produced by *Vibrio cholerae* from patients suffering from choleric diarrhea. Ph.D.Thesis, College of Science, University of Al- Mustansyria.
- 22- Korbayashi, K.; Set, K.; Akasaka, S. and Makino, M. (1990) Detection of toxigenic *Vibrio cholera* O1 using polymerase chain reaction for amplifying the cholera enterotoxin gene. *Kansenshogaku Zasshi. J Jap.Assoc. Infec. Dis.* 64: 1323- 1329.
- 23- Theron, J.; Cilliers, J.; Du Preez, M.; Brozel, V.S. and Nenter, V.S. (2000).Detection of toxigenic *Vibrio cholera* from environmental water samples by an enrichment broth cultivation-pit-stop semi-nested PCR procedure. *Appl. Environ. Microbiol.* 89: 539-546.
- 24- Choopun, N.; Louis, V.; Huq, A.; and Colwell, R.R. (2002). Simple Procedure for Rapid Identification of *Vibrio cholerae*

- from the Aquatic Environment. *Appl. Environ. Microbiol.* 68: 995–998.
- 25- Kaper, J. B.; Morris, J. G. and Levine, M. M. (1995). *Cholera*. *Clin. Microbiol. Rev.* 8:48–86.
- 26- O’Hara, C.M.; Sowers, E.G.; Bopp, C.A.; Duda, S.B. and Strockbine, N.A. (2003). Accuracy of six commercially available systems for identification of members of the family *Vibrionaceae*. *J Clin Microbiol*; 41:5654-9.
- 27- West, P. A. and Colwell, R.R. (1983). Identification and classification of *Vibrionaceae*—an overview, p. 285–341. *In* R. R. Colwell and M. B. Hatem (ed.), *Vibrios in the environment*. John Wiley & Sons, New York, N.Y.
- 28- Chun, J.; Huq, A. and Colwell, R.R. (1999). Analysis of 16S-23S rRNA Intergenic Spacer Regions of *Vibrio cholerae* and *Vibrio mimicus* American Society for Microbiology. 65: 2202–2208.
- 29- Chakraborty, B.; Zaman, K. and Rahman, MD. M. (2008). Rapid Method for Species-Specific Identification and Determination of Toxigenicity of *Vibrio Cholerae* from Natural Aquatic Environment. *S. J. Pharm. Sci.* 1(1&2): 69-75.
- 30- Mousavi, S.L.; Nazarian, S.; Amani, J. and Rahgerdi, A.K. (2008). Rapid Screening of Toxigenic *Vibrio cholerae* O1 Strains from South Iran by PCR-ELISA. *Iran. Biomed. J.* 12 (1): 15-21.
- 31- Koch, W.H.; Payne, W.L.; Wentz, B.A. and Cebula, T.A. (1993) Rapid polymerase chain reaction method for detection of *Vibrio cholera* in foods. *Appl. Environ. Microbiol.* 59: 556-560.